

Date Shipment Needed: _____ Ship To: Patient Prescriber
 Nursing needed; Training needed ► All the supplies including syringes and needles will be dispensed if needed.

CROHN'S DISEASE AND ULCERATIVE COLITIS REFERRAL FORM (A - Si)

PATIENT INFORMATION

Patient Name: _____ DOB: _____ Sex: M F Other: _____ Weight: _____ lbs. kg.
 SSN: _____ Phone: _____ Allergies: _____
 Address: _____ City: _____ State: _____ Zip: _____
 Emergency Contact: _____ Phone: _____ Additional Information Attached

PRESCRIBER INFORMATION

Prescriber: _____ NPI: _____ DEA: _____ State Lic: _____
 Supervising Physician: _____ Practice Name: _____
 Address: _____ City: _____ State: _____ Zip: _____
 Phone: _____ Fax: _____ Key Office Contact: _____ Phone: _____

DIAGNOSIS INFORMATION / MEDICAL ASSESSMENT

Primary Diagnosis: (ICD-10 Code & Description) K50.00 K50.10 K50.80 K50.90 Crohn's Disease K51.9 Ulcerative Colitis Other: _____
 • Has patient been treated *previously* for this condition? Yes No Is patient *currently* on therapy? Yes No Please list medication(s) and treatment duration: _____
 • Will patient stop taking the above medication(s) before starting the new medication? Yes No If yes, how long should patient wait before starting the new medication? _____
 • Other medications patient is currently taking including OTC medications with dosage and direction (or fax medication profile): _____
 • Has patient received a **Quatiferon gold, Tspot or PPD (tuberculosis) Skin Test?** Yes No Date: _____ Results: Negative Positive

INSURANCE INFORMATION

Please attach front and back of patient's insurance card (medical and prescription)

COPAY CARD ENROLLMENT

Please check if enrolling in copay card Copay ID: _____

PRESCRIPTION INFORMATION

STC Standard Protocol will include the following: (1) dispensing ordered med/dose, (2) diluent to mix and/or dilute dose, (3) flushes to flush line and anakit med (epinephrine 0.3 mg IM/0.15 mg IM (for pediatric patients) and diphenhydramine 50 mg/mL) and (4) premeds to take 30 mins before orally (Apap 325 mg, may repeat x1, and diphenhydramine 25 mg, may repeat x1).

Cimzia® 200 mg/mL Prefilled Syringe 200 mg Vial
*Cimzia vial should be prepared and administered by a healthcare professional. Prefilled Syringes will be dispensed unless vial is requested.
 Starter Dose: 400 mg SQ (2 inj. of 200 mg) initially at Week 0, repeat at Weeks 2 and 4 Starter Dose Not Needed QTY: 1 Starter Kit (6 PFS) | Refills: 0
 Maintenance Dose: 400 mg SQ (2 inj. of 200 mg) every 4 weeks QTY: 1 Box (2 x 200 mg) | Refills: _____
 Alternate Dose: _____ QTY: _____ | Refills: _____

Entyvio® 300 mg Vial MD's Office Infusion Home Infusion Supplies Required
 Starter Dose: 300 mg IV at Week 0, Week 2, Week 6 Starter Dose Not Needed QTY: 3 Vials | Refills: 0
 Maintenance Dose: 300 mg IV every 8 weeks QTY: 1 Vial | Refills: _____

Entyvio® 108 mg Pen 108 mg Syringe
 Maintenance Dose: 108 mg SQ once every 2 weeks QTY: 2 Pens/Syringes | Refills: _____
 (beginning after at least 2 IV infusions; administer in place of next scheduled IV dose and then every 2 weeks thereafter)

Humira® CF Starter Package 80 mg/0.8 mL Pen NDC: 0074-0124-03 (See Biosimilar form for alternatives)
 Starter Dose: Two 80 mg SQ inj. Day 1, One 80 mg SQ inj. Day 15 Starter Dose Not Needed QTY: 3 Pens | Refills: 0
 One 80 mg SQ inj. Day 1, One 80 mg SQ inj. Day 2, One 80 mg SQ inj. Day 15 QTY: 3 Pens | Refills: 0

Humira® CF 40 mg/0.4 mL Pen NDC: 0074-0554-02 40 mg/0.4 mL Prefilled Syringe NDC: 0074-0243-02 (See Biosimilar form for alternatives)
 Maintenance Dose: One 40 mg SQ inj. Day 29 & every other week thereafter QTY: 2 | Refills: _____
 Alternate Dose: _____ QTY: _____ | Refills: _____

Omvo® MD's Office Infusion Home Infusion Supplies Required Starter Dose Not Needed
 Starter Dose: 300 mg Vials: 300 mg IV at Weeks 0, 4, and 8 QTY: 1 Vial (28 DS) | Refills: 2
 Maintenance Dose: 100 mg Autoinjector: 200 mg (2 injectors) at Week 12, then every 4 weeks thereafter QTY: 2 Pens (28 DS) | Refills: _____

Remicade® Remicade® 100 mg Vial Inflectra® 100 mg Vial Renflexis® 100 mg Vial Avsola® 100 mg Vial
 MD's Office Infusion Home Infusion Supplies Required Starter Dose Not Needed QTY: _____ | Refills: 0
 Starter Dose: _____ mg IV on Week 0, Week 2, Week 6 QTY: _____ | Refills: _____
 Maintenance Dose: _____ mg IV every _____ weeks QTY: _____ | Refills: _____

Rinvoq®
 Starter Dose: 45 mg Tablet: Once daily x 8 weeks (for Ulcerative Colitis) Starter Dose Not Needed QTY: 28 | Refills: 1
 45 mg Tablet: Once daily x 12 weeks (for Crohn's Disease) QTY: 28 | Refills: 2
 Maintenance Dose: 15 mg Tablet: Once daily QTY: 30 | Refills: _____
 30 mg Tablet: Once daily (alternate maintenance dose for pts. w/ severe or refractory disease) QTY: 30 | Refills: _____

Simponi® SmartJect 100 mg/mL Prefilled Syringe 100 mg/mL *SmartJect will be dispensed unless PFS is requested.
 Starter Dose: 200 mg SQ at Week 0; 100 mg at Week 2; then start maintenance at Week 6 Starter Dose Not Needed QTY: 3 | Refills: 0
 Maintenance Dose: 100 mg SQ every 4 weeks starting at Week 6 QTY: 1 | Refills: _____
 Alternate Dose: _____ QTY: _____ | Refills: _____

Other: _____ QTY: _____ | Refills: _____

Prescriber's Signature: _____ DAW (Dispense as Written) **Date:** _____

Prescriber certifies that this referral form contains an original signature and is signed by the treating prescriber. NO STAMPED SIGNATURES WILL BE ACCEPTED. Where required by law, send electronic prescription or on official state prescription blank. In the event requested agent is not available through AcariaHealth, this prescription shall be forwarded to an eligible pharmacy.

IMPORTANT NOTICE: This message may contain privileged and confidential information and is intended only for the individual named. If you are not the named addressee, you should not disseminate, distribute or copy this fax. Please notify the sender immediately if you have received this document by mistake, then destroy this document. Please direct all verification or notification to AcariaHealth or any of its subsidiaries using the contact information provided on this coversheet.

Date Shipment Needed: _____ Ship To: Patient Prescriber
 Nursing needed; Training needed ► All the supplies including syringes and needles will be dispensed if needed.

CROHN'S DISEASE AND ULCERATIVE COLITIS REFERRAL FORM (Sk - Z)

PATIENT INFORMATION

Patient Name: _____ DOB: _____ Sex: M F Other: _____ Weight: _____ lbs. kg.
 SSN: _____ Phone: _____ Allergies: _____
 Address: _____ City: _____ State: _____ Zip: _____
 Emergency Contact: _____ Phone: _____ Additional Information Attached

PRESCRIBER INFORMATION

Prescriber: _____ NPI: _____ DEA: _____ State Lic: _____
 Supervising Physician: _____ Practice Name: _____
 Address: _____ City: _____ State: _____ Zip: _____
 Phone: _____ Fax: _____ Key Office Contact: _____ Phone: _____

DIAGNOSIS INFORMATION / MEDICAL ASSESSMENT

Primary Diagnosis: (ICD-10 Code & Description) K50.00 K50.10 K50.80 K50.90 Crohn's Disease K51.9 Ulcerative Colitis Other: _____
 • Has patient been treated *previously* for this condition? Yes No Is patient *currently* on therapy? Yes No Please list medication(s) and treatment duration: _____
 • Will patient stop taking the above medication(s) before starting the new medication? Yes No If yes, how long should patient wait before starting the new medication? _____
 • Other medications patient is currently taking including OTC medications with dosage and direction (or fax medication profile): _____
 • Has patient received a **Quatiferon gold, Tspot or PPD (tuberculosis) Skin Test?** Yes No Date: _____ Results: Negative Positive

INSURANCE INFORMATION

Please attach front and back of patient's insurance card (medical and prescription)

COPAY CARD ENROLLMENT

Please check if enrolling in copay card Copay ID: _____

PRESCRIPTION INFORMATION

STC Standard Protocol will include the following: (1) dispensing ordered med/dose, (2) diluent to mix and/or dilute dose, (3) flushes to flush line and anakit med (epinephrine 0.3 mg IM/0.15 mg IM (for pediatric patients) and diphenhydramine 50 mg/mL) and (4) premeds to take 30 mins before orally (Apap 325 mg, may repeat x1, and diphenhydramine 25 mg, may repeat x1).

Skyrizi® MD's Office Infusion Home Infusion Supplies Required
 Starter Dose: **600 mg Vial:** 600 mg IV on Week 0, Week 4, Week 8 Starter Dose Not Needed QTY: _____ Refills: _____
 Maintenance Dose: **360 mg On-Body Injector:** 360 mg SQ on week 12 and every 8 weeks thereafter QTY: _____ Refills: _____
 180 mg On-Body Injector: 180 mg SQ on week 12 and every 8 weeks thereafter QTY: _____ Refills: _____

Stelara® MD's Office Infusion Home Infusion Supplies Required Enroll in Janssen CarePath Program
 Starter Dose: **IV Infusion 130 mg/26 mL (5 mg/mL)** — single-dose vial, weight-based Starter Dose Not Needed
 ≤ 55 kg: IV Infusion 260 mg (2 Vials) once QTY: _____ Refills: _____
 > 55 kg to 85 kg: IV Infusion 390 mg (3 Vials) once QTY: _____ Refills: _____
 > 85 kg: IV Infusion 520 mg (4 Vials) once QTY: _____ Refills: _____
 Maintenance Dose: **90 mg/mL single-dose Prefilled Syringe** QTY: _____ Refills: _____
 Home Injection Dose: SQ inj. 90 mg 8 weeks after first IV dose, every 8 weeks thereafter QTY: _____ Refills: _____

Tremyfa® MD's Office Infusion Home Infusion Supplies Required
 Starter Dose: **200 mg Vial:** 200 mg IV over at least one hour at Week 0, Week 4, Week 8 Starter Dose Not Needed QTY: _____ Refills: _____
 Maintenance Dose: **100 mg Pen** **100 mg Prefilled Syringe** — Administer 100 mg SQ at Week 16, then every 8 weeks thereafter QTY: _____ Refills: _____
 200 mg Pen **200 mg Prefilled Syringe** — Administer 200 mg SQ at Week 12, then every 4 weeks thereafter QTY: _____ Refills: _____

Velsipity® **2 mg Tablet**
 Take 1 Tablet (2 mg) by mouth once daily QTY: _____ Refills: _____

Xeljanz®
 Starter Dose: **10 mg Oral Tablet:** 1 Tablet PO twice daily for 8 weeks Starter Dose Not Needed QTY: _____ Refills: _____
 Other: QTY: _____ Refills: _____
 Maintenance Dose: **5 mg Oral Tablet** **10 mg Oral Tablet** — 1 Tablet PO once daily QTY: _____ Refills: _____
 Other: QTY: _____ Refills: _____

Xeljanz XR®
 Starter Dose: **22 mg Oral Tablet:** 1 Tablet PO once daily for 8 weeks Starter Dose Not Needed QTY: _____ Refills: _____
 Other: QTY: _____ Refills: _____
 Maintenance Dose: **11 mg Oral Tablet** **22 mg Oral Tablet** — 1 Tablet PO once daily QTY: _____ Refills: _____
 Other: QTY: _____ Refills: _____

Zeposia® **Oral Capsules** — Directions: **Days 1-4:** 0.23 mg by mouth once daily; **Days 5-7:** 0.46 mg by mouth once daily;
Day 8 and thereafter: 0.92 mg by mouth once daily
 New Patient: Starter Kit: 7 Day Starter Pack followed by 30 day supply QTY: 1 Kit (37 Capsules) | Refills: 0
 Patients Restarting: 7 Day Titration QTY: 1 Kit (7 Capsules) | Refills: 0
 Maintenance Dose: 0.92 mg by mouth once daily QTY: _____ Refills: _____
 Other: _____ QTY: _____ Refills: _____

Prescriber's Signature: _____ DAW (Dispense as Written) **Date:** _____

Prescriber certifies that this referral form contains an original signature and is signed by the treating prescriber. **NO STAMPED SIGNATURES WILL BE ACCEPTED.** Where required by law, send electronic prescription or on official state prescription blank. In the event requested agent is not available through AcariaHealth, this prescription shall be forwarded to an eligible pharmacy.

IMPORTANT NOTICE: This message may contain privileged and confidential information and is intended only for the individual named. If you are not the named addressee, you should not disseminate, distribute or copy this fax. Please notify the sender immediately if you have received this document by mistake, then destroy this document. Please direct all verification or notification to AcariaHealth or any of its subsidiaries using the contact information provided on this coversheet.