

Date Shipment Needed: \_\_\_\_\_ Ship To:  Patient  Prescriber  
 Nursing needed;  Training needed ► All the supplies including syringes and needles will be dispensed if needed.

**DERMATOLOGY REFERRAL FORM ( A - D )**

**PATIENT INFORMATION**

Patient Name: \_\_\_\_\_ DOB: \_\_\_\_\_ Sex:  M  F  Other: \_\_\_\_\_ Weight: \_\_\_\_\_  lbs.  kg.  
 SSN: \_\_\_\_\_ Phone: \_\_\_\_\_ Allergies: \_\_\_\_\_  
 Address: \_\_\_\_\_ City: \_\_\_\_\_ State: \_\_\_\_\_ Zip: \_\_\_\_\_  
 Emergency Contact: \_\_\_\_\_ Phone: \_\_\_\_\_  Additional Information Attached

**PRESCRIBER INFORMATION**

Prescriber: \_\_\_\_\_ NPI: \_\_\_\_\_ DEA: \_\_\_\_\_ State Lic: \_\_\_\_\_  
 Supervising Physician: \_\_\_\_\_ Practice Name: \_\_\_\_\_  
 Address: \_\_\_\_\_ City: \_\_\_\_\_ State: \_\_\_\_\_ Zip: \_\_\_\_\_  
 Phone: \_\_\_\_\_ Fax: \_\_\_\_\_ Key Office Contact: \_\_\_\_\_ Phone: \_\_\_\_\_

**DIAGNOSIS INFORMATION / MEDICAL ASSESSMENT**

**Primary Diagnosis:**  L20, L20.8, L20.9 Atopic Dermatitis  L28.1 Prurigo nodularis  L40.0 Psoriasis  L40.1; L40.2; L40.3, L40.4, L40.8, L40.54 Psoriatic arthritis  L40.59  
 L50.1 Chronic Idiopathic Urticaria  L73.2 Hidradenitis Suppurativa  Other: \_\_\_\_\_  
 • Location:  Hands  Feet  Face  Scalp  Groin  Nails  Other: \_\_\_\_\_  
 • Severity:  Mild (up to 3% BSA)  Moderate (3-10% BSA)  Severe (greater than 10% BSA), BSA \_\_\_\_\_ %  
 If treated previously for this condition, please indicate which drugs have been tried and failed: \_\_\_\_\_  
 Date range of previous therapy: \_\_\_\_\_  
 • Is patient currently on therapy?  Yes  No Type / medication(s): \_\_\_\_\_  
 • Will patient stop taking the above medication(s) before starting the new medication?  Yes  No, if yes, how long should patient wait before starting the new medication? \_\_\_\_\_  
 • Has patient received a PPD (tuberculosis) Skin Test?  Yes  No Results: \_\_\_\_\_  
 Prior to initiating treatment and periodically during therapy, patient should be evaluated for active tuberculosis and tested for latent infection.

**INSURANCE INFORMATION**

Please attach front and back of patient's insurance card (medical and prescription)

**COPAY CARD ENROLLMENT**

Please check if enrolling in copay card **Copay ID:** \_\_\_\_\_

**PRESCRIPTION INFORMATION**

**Adbry®**  150 mg Prefilled Syringe  
 Starter Dose: 600 mg SQ on Day 1  Starter Dose not needed QTY: #4 Syringes / 14 DS | Refills: 0  
 After 16 wks of treatment  Maintenance Dose: 300 mg SQ every 4 weeks QTY: #2 Syringes / 28 DS | Refills: \_\_\_\_\_  
 Maintenance Dose: 300 mg SQ every other week QTY: #4 Syringes / 28 DS | Refills: \_\_\_\_\_

**Adbry®**  300 mg Pen  
 Starter Dose: 600 mg SQ on Day 1  Starter Dose not needed QTY: #2 Pens / 14 DS | Refills: 0  
 After 16 wks of treatment  Maintenance Dose: 300 mg SQ every 4 weeks QTY: #1 Pens / 28 DS | Refills: \_\_\_\_\_  
 Maintenance Dose: 300 mg SQ every other week QTY: #2 Pens / 28 DS | Refills: \_\_\_\_\_

**Adbry®**  150 mg Prefilled Syringe (Pediatric: 12 years and older)  
 Starter Dose: 300 mg SQ on Day 1  Starter Dose not needed QTY: #2 Syringes / 14 DS | Refills: 0  
 Maintenance Dose: 150 mg SQ every other week QTY: #2 Syringes / 28 DS | Refills: \_\_\_\_\_

**Bimzelx®**  160 mg/mL Pen OR  160 mg/mL Syringe  
 320 mg (given as two 160 mg injections) SQ every 4 weeks for the first 16 weeks QTY: 2 | Refills: 4  
 320 mg (given as two 160 mg injections) SQ every 8 weeks QTY: 2 | Refills: \_\_\_\_\_

**Cibinqo®**  50 mg Tablet  100 mg Tablet  200 mg Tablet  
 1 tablet PO once daily  Other: \_\_\_\_\_ QTY: 1 Month | Refills: \_\_\_\_\_

**Cimzia®**  200 mg Vial  200 mg/mL Prefilled Syringe  
 400 mg/mL SQ every 2 weeks  
 400 mg SQ at Weeks 0, 2, 4, then 200 mg every other week thereafter (patient <=90 kg) QTY: 1 Month | Refills: \_\_\_\_\_

**Cosentyx®**  150 mg/mL Sensoready® Pen  150 mg/mL Prefilled Syringe  300 mg UnoReady Pen  
\*Sensoready® Pen will be dispensed if no preference indicated  
 Starter Dose: 300 mg SQ initially (Weeks 0, 1, 2, 3 and 4) then 300 mg SQ every 4 weeks thereafter (Week 4)  Starter Dose not needed QTY: 5 Weeks | Refills: 0  
 Maintenance Dose: 300 mg SQ every 4 weeks QTY: 1 Month | Refills: \_\_\_\_\_  
 Other: \_\_\_\_\_ QTY: 1 Month | Refills: \_\_\_\_\_

**Dupixent®**  200 mg Pen Autoinjector  200 mg Prefilled Syringe  300 mg Pen Autoinjector  300 mg Prefilled Syringe  
(Dupilumab) \*Pen will be dispensed if no preference indicated for adult dosing. Prefilled Syringe may be used in ages ≥ 6 months. Prefilled Pen is only for use in ages ≥ 2 years.  
**Adults:**  Starter Dose: Inj. 600 mg SQ on Day 1, then 300 mg SQ every 2 weeks starting on Day 15  Starter Dose not needed QTY: QS for Starter | Refills: 0  
 Maintenance Dose: Inj. 300 mg SQ every 2 Weeks QTY: 1 Month | Refills: \_\_\_\_\_  
**Infants & Children:** ≥ 6 mo - < 6 yrs: Initial loading dose not necessary in pediatric patients < 6 yrs.  
 5 to < 15 kg: Dupixent 200 mg SQ every 4 weeks  15 to < 30 kg: Dupixent 300 mg SQ every 4 weeks QTY: 1 Box: 2 Pens/Syringes | Refills: \_\_\_\_\_  
**Children & Adolescents:** ≥ 6 years - ≤ 17 years:  
 15 to < 30 kg: Initial: 600 mg SQ once (administered as two 300 mg injections) QTY: 1 Box: 2 Pens/Syringes | Refills: 0  
 15 to < 30 kg: Maintenance: 300 mg SQ every 4 weeks QTY: 1 Box: 2 Pens/Syringes | Refills: \_\_\_\_\_  
 30 to < 60 kg: Initial: 400 mg SQ once (administered as two 200 mg injections) QTY: 1 Box: 2 Pens/Syringes | Refills: 0  
 30 to < 60 kg: Maintenance: 200 mg SQ every other week QTY: 1 Box: 2 Pens/Syringes | Refills: \_\_\_\_\_  
 ≥ 60 kg: Initial: 600 mg SQ once (administered as two 300 mg injections) QTY: 1 Box: 2 Pens/Syringes | Refills: 0  
 ≥ 60 kg: Maintenance: 300 mg SQ every other week QTY: 1 Box: 2 Pens/Syringes | Refills: \_\_\_\_\_

**Prescriber's Signature:** \_\_\_\_\_  DAW (Dispense as Written) **Date:** \_\_\_\_\_

Prescriber certifies that this referral form contains an original signature and is signed by the treating prescriber. NO STAMPED SIGNATURES WILL BE ACCEPTED. Where required by law, send electronic prescription or on official state prescription blank. In the event requested agent is not available through AcariaHealth, this prescription shall be forwarded to an eligible pharmacy.

**IMPORTANT NOTICE:** This message may contain privileged and confidential information and is intended only for the individual named. If you are not the named addressee, you should not disseminate, distribute or copy this fax. Please notify the sender immediately if you have received this document by mistake, then destroy this document. Please direct all verification or notification to AcariaHealth or any of its subsidiaries using the contact information provided on this coversheet.

Date Shipment Needed: \_\_\_\_\_ Ship To:  Patient  Prescriber  
 Nursing needed;  Training needed ► All the supplies including syringes and needles will be dispensed if needed.

**DERMATOLOGY REFERRAL FORM ( E - Od )**

**PATIENT INFORMATION**

Patient Name: \_\_\_\_\_ DOB: \_\_\_\_\_ Sex:  M  F  Other: \_\_\_\_\_ Weight: \_\_\_\_\_  lbs.  kg.  
 SSN: \_\_\_\_\_ Phone: \_\_\_\_\_ Allergies: \_\_\_\_\_  
 Address: \_\_\_\_\_ City: \_\_\_\_\_ State: \_\_\_\_\_ Zip: \_\_\_\_\_  
 Emergency Contact: \_\_\_\_\_ Phone: \_\_\_\_\_  Additional Information Attached

**PRESCRIBER INFORMATION**

Prescriber: \_\_\_\_\_ NPI: \_\_\_\_\_ DEA: \_\_\_\_\_ State Lic: \_\_\_\_\_  
 Supervising Physician: \_\_\_\_\_ Practice Name: \_\_\_\_\_  
 Address: \_\_\_\_\_ City: \_\_\_\_\_ State: \_\_\_\_\_ Zip: \_\_\_\_\_  
 Phone: \_\_\_\_\_ Fax: \_\_\_\_\_ Key Office Contact: \_\_\_\_\_ Phone: \_\_\_\_\_

**DIAGNOSIS INFORMATION / MEDICAL ASSESSMENT**

**Primary Diagnosis:**  L20, L20.8, L20.9 Atopic Dermatitis  L28.1 Prurigo nodularis  L40.0 Psoriasis  L40.1; L40.2; L40.3, L40.4, L40.8, L40.54 Psoriatic arthritis  L40.59  
 L50.1 Chronic Idiopathic Urticaria  L73.2 Hidradenitis Suppurativa  Other: \_\_\_\_\_  
 • Location:  Hands  Feet  Face  Scalp  Groin  Nails  Other: \_\_\_\_\_  
 • Severity:  Mild (up to 3% BSA)  Moderate (3-10% BSA)  Severe (greater than 10% BSA), BSA \_\_\_\_\_ %  
 If treated previously for this condition, please indicate which drugs have been tried and failed: \_\_\_\_\_  
 Date range of previous therapy: \_\_\_\_\_  
 • Is patient currently on therapy?  Yes  No Type / medication(s): \_\_\_\_\_  
 • Will patient stop taking the above medication(s) before starting the new medication?  Yes  No, if yes, how long should patient wait before starting the new medication? \_\_\_\_\_  
 • Has patient received a PPD (tuberculosis) Skin Test?  Yes  No Results: \_\_\_\_\_  
 Prior to initiating treatment and periodically during therapy, patient should be evaluated for active tuberculosis and tested for latent infection.

**INSURANCE INFORMATION**

Please attach front and back of patient's insurance card (medical and prescription)

**COPY CARD ENROLLMENT**

Please check if enrolling in copay card **Copay ID:** \_\_\_\_\_

**PRESCRIPTION INFORMATION**

<p><b>Enbrel®</b> <input type="checkbox"/> 50 mg/mL SureClick (Autoinjector) <input type="checkbox"/> 50 mg Prefilled Syringe <input type="checkbox"/> Mini 50 mg Cartridge *SureClick will be dispensed if no preference indicated  <input type="checkbox"/> Starter Dose: 50 mg SQ twice weekly (72 - 96 hours apart) for 3 months <input type="checkbox"/> Starter Dose not needed  <input type="checkbox"/> Maintenance Dose: 50 mg SQ weekly <input type="checkbox"/> Other: _____</p> <p><b>Enbrel®</b> <input type="checkbox"/> 25 mg/0.5 mL Prefilled Syringe <input type="checkbox"/> 25 mg Single-Use Vial *Prefilled Syringe will be dispensed if no preference indicated  <input type="checkbox"/> 25 mg SQ twice weekly (72 - 96 hours apart) <input type="checkbox"/> Other: _____</p> <p><b>Eriedge®</b> <input type="checkbox"/> 150 mg Capsules Take 1 capsule orally once daily</p> <p><b>Humira®</b> <input type="checkbox"/> CF Pen Psoriasis Starter Kit NDC: 0074-1539-03 <input type="checkbox"/> CF 40 mg/0.4 mL Prefilled Syringe NDC: 0074-0243-02  <small>*Pen Starter Kit will be dispensed if no preference indicated</small>  <input type="checkbox"/> Starter Dose <input type="checkbox"/> One 80 mg SQ inj. Day 1, one 40 mg SQ inj. Day 8, one 40 mg SQ inj. Day 22 (OR) <input type="checkbox"/> Starter Dose not needed                  for Psoriasis: <input type="checkbox"/> Two 40 mg SQ inj. Day 1, one 40 mg SQ inj. Day 8, one 40 mg SQ inj. Day 22</p> <p><b>Humira®</b> <input type="checkbox"/> CF 40 mg/0.4 mL Pen NDC: 0074-0554-02 <input type="checkbox"/> CF 40 mg/0.4 mL Syringe NDC: 0074-0243-02  <small>*Pen will be dispensed if no preference indicated</small>  <input type="checkbox"/> Maintenance Dose for Psoriasis: 40 mg SQ once every other week</p> <p><b>Humira®</b> <input type="checkbox"/> Starter Pkg CF 80 mg/0.8 mL Pen NDC: 0074-0124-03 <input type="checkbox"/> CF 40 mg/0.4 mL Prefilled Syringe NDC: 0074-0243-02  <small>*Pen will be dispensed if no preference indicated</small>  <input type="checkbox"/> Starter Dose for <input type="checkbox"/> Inj. 160 mg SQ Day 1, then 80 mg SQ Day 15 (OR) <input type="checkbox"/> Starter Dose not needed                  Hidradenitis Suppurativa: <input type="checkbox"/> Inj. 80 mg SQ Day 1, and 80 mg SQ Day 2, then 80 mg SQ Day 15</p> <p><b>Humira®</b> <input type="checkbox"/> CF 40 mg/0.4 mL Pen NDC: 0074-0554-02 <input type="checkbox"/> CF 40 mg/0.4 mL Syringe NDC: 0074-0243-02  <small>*Pen will be dispensed if no preference indicated</small>  <input type="checkbox"/> Maintenance Dose for Hidradenitis Suppurativa: 40 mg SQ Day 29 and every week thereafter  <input type="checkbox"/> Other: _____</p> <p><b>Ilumya®</b> <input type="checkbox"/> 100 mg/mL Prefilled Syringes  <input type="checkbox"/> Starter Dose: 100 mg SQ on Week 0 and Week 4 <input type="checkbox"/> Starter Dose not needed  <input type="checkbox"/> Maintenance Dose: 100 mg SQ every 12 weeks (starting at Week 4)</p> <p><b>Kevzara®</b> <input type="checkbox"/> 200 mg/1.14 mL Pen Autoinjector <input type="checkbox"/> 200 mg/1.14 mL Prefilled Syringe *Pens will be dispensed if no preference is indicated                  (Sarilumab) <input type="checkbox"/> 200 mg subcutaneously every 2 Weeks</p> <p><b>Nemluvio®</b> <input type="checkbox"/> 30 mg Pen - The recommended subcutaneous dosage of NEMLUVIO for adult patients weighing less than 90 kg is an initial dose of 60 mg (two 30 mg injections), followed by 30 mg given every 4 weeks (Q4W)                  Adult Patients Weighing less than 90 kg: <input type="checkbox"/> Inject 60 mg SQ once for initial dose  <input type="checkbox"/> Inject 30 mg SQ every 4 weeks</p> <p><b>Nemluvio®</b> <input type="checkbox"/> 60 mg Pen - The recommended subcutaneous dosage of NEMLUVIO for adult patients weighing 90 kg or more is an initial dose of 60 mg (two 30 mg injections), followed by 60 mg given every 4 weeks (Q4W)                  Adult Patients Weighing 90 kg or more: <input type="checkbox"/> Inject 60 mg SQ once for initial dose  <input type="checkbox"/> Inject 60 mg SQ every 4 weeks</p> <p><b>Odanzo®</b> <input type="checkbox"/> 200 mg Capsule PO Once Daily</p>	<p><input type="checkbox"/> Enroll in Enliven® Program                  QTY: 1 Month   Refills: 2                  QTY: 1 Month   Refills: _____</p> <p>QTY: 1 Month   Refills: _____                  QTY: 28 Capsules   Refills: _____</p> <p><input type="checkbox"/> Enroll in Humira Complete Program                  QTY: 3 Pens   Refills: 0                  QTY: 4 Syringes   Refills: 0</p> <p>QTY: 1 Month   Refills: _____                  QTY: 1 Month   Refills: 0                  QTY: 1 Month   Refills: 0</p> <p>QTY: 1 Month   Refills: _____                  QTY: _____   Refills: _____</p> <p>QTY: 1 Month (1 PFS)   Refills: 0                  QTY: 1 Syringe   Refills: _____</p> <p>QTY: 1 Box (2)   Refills: _____</p> <p>QTY: #2 Pens/28 DS   Refills: 0                  QTY: #1 Pens/28 DS   Refills: _____</p> <p>QTY: #2 Pens/28 DS   Refills: 0                  QTY: #2 Pens/28 DS   Refills: _____</p> <p>QTY: 30 CAPS   Refills: _____</p>
---	---

**Prescriber's Signature:** \_\_\_\_\_  DAW (Dispense as Written) **Date:** \_\_\_\_\_

Prescriber certifies that this referral form contains an original signature and is signed by the treating prescriber. NO STAMPED SIGNATURES WILL BE ACCEPTED. Where required by law, send electronic prescription or on official state prescription blank. In the event requested agent is not available through AcariaHealth, this prescription shall be forwarded to an eligible pharmacy.

**IMPORTANT NOTICE:** This message may contain privileged and confidential information and is intended only for the individual named. If you are not the named addressee, you should not disseminate, distribute or copy this fax. Please notify the sender immediately if you have received this document by mistake, then destroy this document. Please direct all verification or notification to AcariaHealth or any of its subsidiaries using the contact information provided on this coversheet.

Date Shipment Needed: \_\_\_\_\_ Ship To:  Patient  Prescriber  
 Nursing needed;  Training needed ▶ All the supplies including syringes and needles will be dispensed if needed.

**DERMATOLOGY REFERRAL FORM ( Ot - So )**

**PATIENT INFORMATION**

Patient Name: \_\_\_\_\_ DOB: \_\_\_\_\_ Sex:  M  F  Other: \_\_\_\_\_ Weight: \_\_\_\_\_  lbs.  kg.  
 SSN: \_\_\_\_\_ Phone: \_\_\_\_\_ Allergies: \_\_\_\_\_  
 Address: \_\_\_\_\_ City: \_\_\_\_\_ State: \_\_\_\_\_ Zip: \_\_\_\_\_  
 Emergency Contact: \_\_\_\_\_ Phone: \_\_\_\_\_  Additional Information Attached

**PRESCRIBER INFORMATION**

Prescriber: \_\_\_\_\_ NPI: \_\_\_\_\_ DEA: \_\_\_\_\_ State Lic: \_\_\_\_\_  
 Supervising Physician: \_\_\_\_\_ Practice Name: \_\_\_\_\_  
 Address: \_\_\_\_\_ City: \_\_\_\_\_ State: \_\_\_\_\_ Zip: \_\_\_\_\_  
 Phone: \_\_\_\_\_ Fax: \_\_\_\_\_ Key Office Contact: \_\_\_\_\_ Phone: \_\_\_\_\_

**DIAGNOSIS INFORMATION / MEDICAL ASSESSMENT**

**Primary Diagnosis:**  L20, L20.8, L20.9 Atopic Dermatitis  L28.1 Prurigo nodularis  L40.0 Psoriasis  L40.1; L40.2; L40.3, L40.4, L40.8, L40.54 Psoriatic arthritis  L40.59  
 L50.1 Chronic Idiopathic Urticaria  L73.2 Hidradenitis Suppurativa  Other: \_\_\_\_\_  
 • Location:  Hands  Feet  Face  Scalp  Groin  Nails  Other: \_\_\_\_\_  
 • Severity:  Mild (up to 3% BSA)  Moderate (3-10% BSA)  Severe (greater than 10% BSA), BSA \_\_\_\_\_ %  
 If treated previously for this condition, please indicate which drugs have been tried and failed: \_\_\_\_\_  
 Date range of previous therapy: \_\_\_\_\_  
 • Is patient currently on therapy?  Yes  No Type / medication(s): \_\_\_\_\_  
 • Will patient stop taking the above medication(s) before starting the new medication?  Yes  No, if yes, how long should patient wait before starting the new medication? \_\_\_\_\_  
 • Has patient received a PPD (tuberculosis) Skin Test?  Yes  No Results: \_\_\_\_\_  
 Prior to initiating treatment and periodically during therapy, patient should be evaluated for active tuberculosis and tested for latent infection.

**INSURANCE INFORMATION**

Please attach front and back of patient's insurance card (medical and prescription)

**COPAY CARD ENROLLMENT**

Please check if enrolling in copay card **Copay ID:** \_\_\_\_\_

**PRESCRIPTION INFORMATION**

**Otezla® Tablets** Plaque Psoriasis - moderate to severe; **Note:** Initial dose titration is intended to reduce GI symptoms

— **Adults:**  
 Titration **DAY 1:** 10 mg in morning; **DAY 2:** 10 mg in morning & 10 mg in evening; **DAY 3:** 10 mg in morning & 20 mg in evening; QTY: \_\_\_\_\_ 1 Month | Refills: 0  
 Dose: **DAY 4:** 20 mg in morning & 20 mg in evening; **DAY 5:** 20 mg in morning & 30 mg in evening; **DAY 6 & thereafter:** 30 mg twice daily  
 Maintenance Dose: 30 mg twice daily QTY: 60 TABS (30 mg) | Refills: \_\_\_\_\_  
 Other: \_\_\_\_\_ QTY: \_\_\_\_\_ | Refills: \_\_\_\_\_

— **Children & Adolescents:** ≥ 6 years weighing 20 to < 50 kg:  
 Otezla® 28 Day Treatment Initiation Pack: **DAY 1:** Oral: 10 mg once daily in the morning; **DAY 2:** Oral: 10 mg twice daily; QTY: \_\_\_\_\_ # 1 | Refills: 0  
**DAY 3:** Oral: 10 mg in the morning & 20 mg in the evening; **DAY 4 & thereafter:** Oral: 20 mg twice daily  
 Otezla® 20 mg Tablet: Take 20 mg PO twice daily QTY: 60 TABS | Refills: \_\_\_\_\_

— **Children & Adolescents:** ≥ 6 years weighing ≥ 50 kg:  
 Otezla® 28 Day Treatment Initiation Pack: **DAY 1:** Oral: 10 mg once daily in the morning; **DAY 2:** Oral: 10 mg twice daily; QTY: \_\_\_\_\_ # 1 | Refills: 0  
**DAY 3:** Oral: 10 mg in the morning & 20 mg in the evening; **DAY 4:** Oral: 20 mg twice daily;  
**DAY 5:** Oral: 20 mg in the morning & 30 mg in the evening; **DAY 6 & thereafter:** Oral: 30 mg twice daily  
 Otezla® 30 mg Tablet: Take 30 mg PO twice daily QTY: 60 TABS | Refills: \_\_\_\_\_

Remicade® 100 mg Vial  Inflectra® 100 mg Powder Vial  Renflexis® 100 mg Powder Vial  Avsola® 100 mg Powder Vial  
 MD's Office Infusion  Home Infusion Supplies Required  Starter Dose not needed  
 Starter Dose: \_\_\_\_\_ mg IV on Week 0, Week 2, Week 6, then QTY: QS 3 Infusions | Refills: 0  
 Maintenance Dose: \_\_\_\_\_ mg IV every \_\_\_\_\_ weeks QTY: QS 1 Infusion | Refills: \_\_\_\_\_

**Rinvoq®**  15 mg Tablet  30 mg Tablet Take 1 tablet PO once daily QTY: 1 Month | Refills: \_\_\_\_\_

**Siliq®**  210 mg / 1.5 mL Prefilled Syringe (2 pack)  Starter Dose not needed  Enroll in REMS Program  
 Starter Dose for Plaque Psoriasis: 210 mg SQ at Weeks 0, 1 and 2, followed by maintenance dose QTY: 1 Box (2 PFS) | Refills: 0  
 Maintenance Dose for Plaque Psoriasis: 210 mg SQ once every two weeks (Starting at Week 2) QTY: 1 Box (2 PFS) | Refills: \_\_\_\_\_

**Simponi®**  Aria 50 mg / 4 mL Patient Weight (kg): \_\_\_\_\_  Starter Dose not needed  Enroll in SimponiOne® Program  
 Starter Dose: 2 mg / kg IV at Weeks 0 and 4 QTY: 1 Month | Refills: 0  
 Maintenance Dose: 2 mg / kg IV every 8 weeks QTY: QS for 8 Weeks | Refills: \_\_\_\_\_

**Simponi®**  SmartJect 50 mg / 0.5 mL  50 mg / 0.5 mL Prefilled Syringe \*Pens will be dispensed if no preference is indicated  
 50 mg SQ every month QTY: 1 Month | Refills: \_\_\_\_\_  
 Other: \_\_\_\_\_ QTY: \_\_\_\_\_ | Refills: \_\_\_\_\_

**Skyrizi®**  150 mg / mL Pen Autoinjector  150 mg / mL Prefilled Syringe \*Pens will be dispensed if no preference is indicated  
 Starter Dose: 150 mg SQ at Week 0 and 4  Starter Dose not needed QTY: 1 | Refills: 1  
 Maintenance Dose: 150 mg SQ every 12 weeks QTY: 1 | Refills: \_\_\_\_\_

**Sotyku®**  6 mg PO once daily QTY: \_\_\_\_\_ | Refills: \_\_\_\_\_

**Prescriber's Signature:** \_\_\_\_\_  DAW (Dispense as Written) **Date:** \_\_\_\_\_

Prescriber certifies that this referral form contains an original signature and is signed by the treating prescriber. NO STAMPED SIGNATURES WILL BE ACCEPTED. Where required by law, send electronic prescription or official state prescription blank. In the event requested agent is not available through AcariaHealth, this prescription shall be forwarded to an eligible pharmacy.

**IMPORTANT NOTICE:** This message may contain privileged and confidential information and is intended only for the individual named. If you are not the named addressee, you should not disseminate, distribute or copy this fax. Please notify the sender immediately if you have received this document by mistake, then destroy this document. Please direct all verification or notification to AcariaHealth or any of its subsidiaries using the contact information provided on this coversheet.

Date Shipment Needed: \_\_\_\_\_ Ship To:  Patient  Prescriber

Nursing needed;  Training needed ▶ All the supplies including syringes and needles will be dispensed if needed.

## DERMATOLOGY REFERRAL FORM ( St - Z )

PATIENT INFORMATION					
Patient Name:		DOB:	Sex: <input type="checkbox"/> M <input type="checkbox"/> F <input type="checkbox"/> Other:	Weight:	<input type="checkbox"/> lbs. <input type="checkbox"/> kg.
SSN:	Phone:	Allergies:			
Address:			City:	State:	Zip:
Emergency Contact:		Phone:	<input type="checkbox"/> Additional Information Attached		

PRESCRIBER INFORMATION					
Prescriber:		NPI:	DEA:	State Lic:	
Supervising Physician:			Practice Name:		
Address:			City:	State:	Zip:
Phone:	Fax:	Key Office Contact:		Phone:	

DIAGNOSIS INFORMATION / MEDICAL ASSESSMENT	
<p><b>Primary Diagnosis:</b> <input type="checkbox"/> L20, L20.8, L20.9 Atopic Dermatitis <input type="checkbox"/> L28.1 Prurigo nodularis <input type="checkbox"/> L40.0 Psoriasis <input type="checkbox"/> L40.1; L40.2; L40.3, L40.4, L40.8, L40.54 Psoriatic arthritis <input type="checkbox"/> L40.59 <input type="checkbox"/> L50.1 Chronic Idiopathic Urticaria <input type="checkbox"/> L73.2 Hidradenitis Suppurativa <input type="checkbox"/> Other: _____</p> <p>• Location: <input type="checkbox"/> Hands <input type="checkbox"/> Feet <input type="checkbox"/> Face <input type="checkbox"/> Scalp <input type="checkbox"/> Groin <input type="checkbox"/> Nails <input type="checkbox"/> Other: _____</p> <p>• Severity: <input type="checkbox"/> Mild (up to 3% BSA) <input type="checkbox"/> Moderate (3-10% BSA) <input type="checkbox"/> Severe (greater than 10% BSA), BSA _____%</p> <p>If treated previously for this condition, please indicate which drugs have been tried and failed: _____</p> <p>Date range of previous therapy: _____</p> <p>• Is patient currently on therapy? <input type="checkbox"/> Yes <input type="checkbox"/> No Type / medication(s): _____</p> <p>• Will patient stop taking the above medication(s) before starting the new medication? <input type="checkbox"/> Yes <input type="checkbox"/> No, if yes, how long should patient wait before starting the new medication? _____</p> <p>• Has patient received a PPD (tuberculosis) Skin Test? <input type="checkbox"/> Yes <input type="checkbox"/> No Results: _____</p> <p>Prior to initiating treatment and periodically during therapy, patient should be evaluated for active tuberculosis and tested for latent infection.</p>	

INSURANCE INFORMATION	
<input type="checkbox"/> Please attach front and back of patient's insurance card (medical and prescription)	

COPAY CARD ENROLLMENT	
<input type="checkbox"/> Please check if enrolling in copay card	Copay ID: _____

PRESCRIPTION INFORMATION	
<p><b>Stelara®</b> <input type="checkbox"/> Prefilled Syringe <input type="checkbox"/> Vial <input type="checkbox"/> MD's Office Infusion <input type="checkbox"/> Home Infusion Supplies Required <i>*Prefilled Syringe will be dispensed if preference is not indicated</i></p> <p><input type="checkbox"/> ≤ 100 kg Starter Dose: 45 mg SQ initially (Week 0), then 45 mg SQ after 4 weeks of initial dose (Week 4)</p> <p><input type="checkbox"/> ≤ 100 kg Maintenance Dose: 45 mg SQ every 12 weeks <input type="checkbox"/> Other: _____</p> <p><input type="checkbox"/> &gt; 100 kg Starter Dose: 90 mg SQ initially (Week 0), then 90 mg SQ after 4 weeks of initial dose (Week 4)</p> <p><input type="checkbox"/> &gt; 100 kg Maintenance Dose: 90 mg SQ every 12 weeks <input type="checkbox"/> Other: _____</p> <p><b>Taltz®</b> <input type="checkbox"/> 80 mg/mL Autoinjector <input type="checkbox"/> 80 mg/mL Prefilled Syringe <i>*Pen will be dispensed if no preference is indicated</i> <input type="checkbox"/> Starter Dose not needed</p> <p><input type="checkbox"/> Starter Dose for Plaque Psoriasis: 160 mg (two 80 mg inj.) at Week 0, then 80 mg at Week 2, 4, 6, 8, 10, 12</p> <p><input type="checkbox"/> Maintenance Dose for Plaque Psoriasis: 80 mg every 4 weeks</p> <p><input type="checkbox"/> Starter Dose for Psoriatic Arthritis: 160 mg (two 80 mg inj.) at Week 0</p> <p><input type="checkbox"/> Maintenance Dose for Psoriatic Arthritis: 80 mg every 4 weeks</p> <p><input type="checkbox"/> Other: _____</p> <p><b>Tremfya®</b> <input type="checkbox"/> 100 mg/mL Pen Autoinjector <input type="checkbox"/> 100 mg/ml Prefilled Syringe <i>*Pens will be dispensed if no preference is indicated</i></p> <p><input type="checkbox"/> Starter Dose: 100 mg SQ at Week 0, 4, and every 8 weeks thereafter <input type="checkbox"/> Starter Dose not needed</p> <p><input type="checkbox"/> Maintenance Dose: 100 mg SQ every 8 weeks (starting at Week 4)</p> <p><b>Xeljanz®</b> <input type="checkbox"/> 5 mg Tablet <input type="checkbox"/> 10 mg Tablet: 1 tablet PO twice daily</p> <p><input type="checkbox"/> 11 mg ER Tablet: 1 tablet PO daily</p> <p><b>Xolair®</b> <input type="checkbox"/> 150 mg Prefilled Syringe <input type="checkbox"/> 150 mg Vial</p> <p><input type="checkbox"/> 150 mg SQ every 4 weeks <input type="checkbox"/> 300 mg SQ every 4 weeks</p>	<p><input type="checkbox"/> Enroll in Janssen CarePath Program</p> <p>QTY: <u>1 x 45 mg</u>   Refills: <u>1</u></p> <p>QTY: <u>1 x 45 mg</u>   Refills: _____</p> <p>QTY: <u>1 x 90 mg</u>   Refills: <u>1</u></p> <p>QTY: <u>1 x 90 mg</u>   Refills: _____</p> <p>QTY: <u>8</u>   Refills: <u>0</u></p> <p>QTY: <u>1</u>   Refills: _____</p> <p>QTY: <u>2</u>   Refills: <u>0</u></p> <p>QTY: <u>1</u>   Refills: _____</p> <p>QTY: _____   Refills: _____</p> <p>QTY: <u>1</u>   Refills: <u>0</u></p> <p>QTY: <u>1</u>   Refills: _____</p> <p>QTY: <u>1 Month</u>   Refills: _____</p> <p>QTY: <u>1 Month</u>   Refills: _____</p> <p>QTY: <u>28 Day Supply</u>   Refills: _____</p>

**Prescriber's Signature:** \_\_\_\_\_  DAW (Dispense as Written) **Date:** \_\_\_\_\_

Prescriber certifies that this referral form contains an original signature and is signed by the treating prescriber. NO STAMPED SIGNATURES WILL BE ACCEPTED. Where required by law, send electronic prescription or on official state prescription blank. In the event requested agent is not available through AcariaHealth, this prescription shall be forwarded to an eligible pharmacy.

**IMPORTANT NOTICE:** This message may contain privileged and confidential information and is intended only for the individual named. If you are not the named addressee, you should not disseminate, distribute or copy this fax. Please notify the sender immediately if you have received this document by mistake, then destroy this document. Please direct all verification or notification to AcariaHealth or any of its subsidiaries using the contact information provided on this coversheet.