

Date Shipment Needed: _____ Ship To: Patient Prescriber
 Nursing needed; Training needed ▶ All the supplies including syringes and needles will be dispensed if needed.

RHEUMATOLOGY NON-IV REFERRAL FORM A-OL

PATIENT INFORMATION

Patient Name: _____ DOB: _____ Sex: M F Other: _____ Weight: _____ lbs. kg.
 SSN: _____ Phone: _____ Allergies: _____
 Address: _____ City: _____ State: _____ Zip: _____
 Emergency Contact: _____ Phone: _____ Additional Information Attached

PRESCRIBER INFORMATION

Prescriber: _____ NPI: _____ DEA: _____ State Lic: _____
 Supervising Physician: _____ Practice Name: _____
 Address: _____ City: _____ State: _____ Zip: _____
 Phone: _____ Fax: _____ Key Office Contact: _____ Phone: _____

DIAGNOSIS INFORMATION / MEDICAL ASSESSMENT

Primary Diagnosis: M06.9 Rheumatoid Arthritis L40.54; L40.59 Psoriatic Arthritis M08.00 Unspecified Juvenile Rheumatoid Arthritis M08.3 Juvenile Rheumatoid Polyarthritis (Seronegative)
 M08.20 Juvenile Idiopathic Arthritis M45.9 Ankylosing Spondylitis M33.20 Polymyositis M81.0 Osteoporosis M15.0; M15.9 Osteoarthritis Other: _____
 • Has patient been treated *previously* for this condition? Yes No Is patient *currently* on therapy? Yes No Please list medication(s) and treatment duration: _____
 • Will patient stop taking the above medication(s) before starting the new medication? Yes No If yes, how long should patient wait before starting the new medication? _____
 • Other medications patient is currently taking including OTC medications with dosage and direction (or fax medication profile): _____
 • Has patient received a **Quatiferon gold, Tspot or PPD (tuberculosis) Skin Test?** Yes No Date: _____ Results: Negative Positive
Prior to initiating treatment and periodically during therapy, patient should be evaluated for active tuberculosis and tested for latent infection

INSURANCE INFORMATION

Please attach front and back of patient's insurance card (medical and prescription)

COPY CARD ENROLLMENT

Please check if enrolling in copay card **Copay ID:** _____

PRESCRIPTION INFORMATION

Actemra® 162 mg Pen Prefilled Syringe
 <100 kg: 162 mg SQ once every other week QTY: 2 Pens / Syringes | Refills: _____
 ≥100 kg: 162 mg SQ once every other week QTY: 4 Pens / Syringes | Refills: _____

Cimzia® 200 mg/mL Prefilled Syringe **Cimzia® 200 mg Vial** Starter Dose Not Needed
**Cimzia vial should be prepared and administered by a healthcare professional. Prefilled Syringe will be dispensed unless vial is requested.*
 Starter Dose: 400 mg SQ (2 inj. of 200 mg) initially (Week 0), repeat at Weeks 2 and 4
 Maintenance Dose: 400 mg SQ (2 inj. of 200 mg) every 4 weeks 200 mg SQ every 2 weeks
 Alternate Dose: _____
 Enroll in Enroll in Cimplicity™ Program
 QTY: 1 Starter Kit (6 PFS) | Refills: 0
 QTY: 1 Box (2 INJ.) | Refills: _____
 QTY: _____ | Refills: _____

Cosentyx® 150 mg/mL Prefilled Syringe **Cosentyx® 150 mg/mL Sensoready Pen** **Cosentyx® 300 mg/2 mL UnoReady Pen**
 Starter Dose: 150 mg SQ at Weeks 0, 1, 2, 3 300 mg SQ at Weeks 0, 1, 2, 3 Starter Dose Not Needed
 Maintenance Dose: 150 mg SQ every 4 weeks (starting at Week 4) 300 mg SQ every 4 weeks (starting at Week 4)
 QTY: QS 28 Day Supply | Refills: 0
 QTY: QS 28 Day Supply | Refills: _____

Enbrel® 50 mg/mL Sureclick (Autoinjector) **Enbrel® 50 mg/mL Prefilled Syringe** **Not to be used in pediatric weighing less than 63 kg (138 lb.)*
 50 mg SQ weekly
 Alternate Dose: _____
 Enroll in Enliven® Program
 QTY: 4 | Refills: _____
 QTY: _____ | Refills: _____

Enbrel® 25 mg/0.5 mL Prefilled Syringe
 25 mg SQ twice weekly (72-96 hours apart)
 Alternate Dose: _____
 QTY: 8 | Refills: _____
 QTY: _____ | Refills: _____

Humira® 40 mg/0.4 mL Pen CF **Humira® 40 mg/0.4 mL Prefilled Syringe CF**
 inject 40 mg SQ every other week inject 40 mg SQ every week
 Enroll in Humira Complete Program
 QTY: _____ | Refills: _____

Humira® 10 mg/0.2 mL Prefilled Syringe CF **Humira® 20 mg/0.4 mL Prefilled Syringe CF**
 inject 10 mg SQ every other week inject 20 mg SQ every other week
 QTY: _____ | Refills: _____
 QTY: _____ | Refills: _____

Ilaris® 150 mg SDV Inj. _____ mg SQ every _____ Weeks
 QTY: _____ | Refills: _____

Ilaris® 150 mg/mL Single Dose Vial
 Inject 4 mg/kg _____ mg SQ every 4 weeks (300 mg/dose maximum)
 QTY: _____ | Refills: _____

Kevzara® Inj. Single Prefilled Syringe **Kevzara® Inj. Single Prefilled Pen**
 150 mg/1.14 mL 200 mg/1.14 mL 1 SQ inj. every 2 weeks
 QTY: _____ | Refills: _____

Olumiant® 2 mg Tablet **1 mg Tablet**
 take 2 mg tablet PO once daily
 take 1 mg tablet PO once daily
 QTY: _____ | Refills: _____
 QTY: _____ | Refills: _____

Prescriber's Signature: _____ DAW (Dispense as Written) **Date:** _____

Prescriber certifies that this referral form contains an original signature and is signed by the treating prescriber. NO STAMPED SIGNATURES WILL BE ACCEPTED. Where required by law, send electronic prescription or on official state prescription blank. In the event requested agent is not available through AcariaHealth, this prescription shall be forwarded to an eligible pharmacy.

IMPORTANT NOTICE: This message may contain privileged and confidential information and is intended only for the individual named. If you are not the named addressee, you should not disseminate, distribute or copy this fax. Please notify the sender immediately if you have received this document by mistake, then destroy this document. Please direct all verification or notification to AcariaHealth or any of its subsidiaries using the contact information provided on this coversheet.

Date Shipment Needed: _____ Ship To: Patient Prescriber
 Nursing needed; Training needed ► All the supplies including syringes and needles will be dispensed if needed.

RHEUMATOLOGY NON-IV REFERRAL FORM OM-Z

PATIENT INFORMATION

Patient Name: _____ DOB: _____ Sex: M F Other: _____ Weight: _____ lbs. kg.
 SSN: _____ Phone: _____ Allergies: _____
 Address: _____ City: _____ State: _____ Zip: _____
 Emergency Contact: _____ Phone: _____ Additional Information Attached

PRESCRIBER INFORMATION

Prescriber: _____ NPI: _____ DEA: _____ State Lic: _____
 Supervising Physician: _____ Practice Name: _____
 Address: _____ City: _____ State: _____ Zip: _____
 Phone: _____ Fax: _____ Key Office Contact: _____ Phone: _____

DIAGNOSIS INFORMATION / MEDICAL ASSESSMENT

Primary Diagnosis: M06.9 Rheumatoid Arthritis L40.54; L40.59 Psoriatic Arthritis M08.00 Unspecified Juvenile Rheumatoid Arthritis M08.3 Juvenile Rheumatoid Polyarthritis (Seronegative)
 M08.20 Juvenile Idiopathic Arthritis M45.9 Ankylosing Spondylitis M33.20 Polymyositis M81.0 Osteoporosis M15.0; M15.9 Osteoarthritis Other: _____
 • Has patient been treated *previously* for this condition? Yes No Is patient *currently* on therapy? Yes No Please list medication(s) and treatment duration: _____
 • Will patient stop taking the above medication(s) before starting the new medication? Yes No If yes, how long should patient wait before starting the new medication? _____
 • Other medications patient is currently taking including OTC medications with dosage and direction (or fax medication profile): _____
 • Has patient received a **Quatiferon gold, Tspot or PPD (tuberculosis) Skin Test?** Yes No Date: _____ Results: Negative Positive
Prior to initiating treatment and periodically during therapy, patient should be evaluated for active tuberculosis and tested for latent infection

INSURANCE INFORMATION

Please attach front and back of patient's insurance card (medical and prescription)

COPY CARD ENROLLMENT

Please check if enrolling in copay card **Copay ID:** _____

PRESCRIPTION INFORMATION

Otezla® Tablet Five (5) day titration period: **Day 1:** 10 mg, **Day 2:** 10 mg BID, **Day 3:** 10 mg in AM then 20 mg in PM, **Day 4:** 20 mg BID, **Day 5:** 20 mg in AM then 30 mg in PM QTY: _____ 1 Kit | Refills: 0
 After five (5) day titration period, 30 mg BID QTY: _____ 60 Tablets | Refills: _____

Orencia® 125 mg Prefilled Syringe **Orencia® 125 mg ClickJect Autoinjector**
 Starter Dose: One dose of IV infusion (per body weight) IV Starter Dose Not Needed
 <60 kg: 500 mg IV x 1 dose
 60-100 kg: 750 mg IV x 1 dose
 >100 kg: 1000 mg IV x 1 dose
 Maintenance Dose: 150 mg SQ every week
 Enroll in Orencia OnCall Program
 QTY: 2 x 250 mg Vial | Refills: 0
 QTY: 3 x 250 mg Vial | Refills: 0
 QTY: 4 x 250 mg Vial | Refills: 0
 QTY: 4 PFS / Pens | Refills: _____

Rinvoq® 15 mg Oral Tablet
 Take one tablet orally once daily with or without food QTY: _____ 30 | Refills: _____

Siliq® 210 mg / 1.5 mL Prefilled Syringe
 210 mg SQ at Weeks 0, 1, 2 QTY: _____ 2 Pens | Refills: 0
 210 mg SQ every 2 weeks (starting at week 2) QTY: _____ 2 Pens | Refills: _____

Skyrizi® 150 mg/mL Pen **Skyrizi® 150 mg/mL Prefilled Syringe**
 Inj. 150 mg SQ at Week 0 QTY: _____ 1 | Refills: 0
 Inj. 150 mg SQ every 12 weeks (starting at Week 4) QTY: _____ 1 | Refills: _____

Simponi® 50 mg / 0.5 mL SmartJect (Autoinjector) **Simponi® 50 mg / 0.5 mL Prefilled Syringe**
 50 mg SQ every month
 Alternate Dose: _____
 Enroll in SimponiOne Program
 QTY: _____ | Refills: _____
 QTY: _____ | Refills: _____

Taltz® 80 mg/mL **Pen** **Prefilled Syringe** Starter Dose Not Needed
 Starter Dose: inject 160 mg once SQ on Week 0 QTY: _____ 2 | Refills: _____
 Maintenance Dose: inject 80 mg once SQ every 4 weeks QTY: _____ 1 Pen / 1 Syringe | Refills: _____

Tremfya® 100 mg/mL **Pen** **Prefilled Syringe** Starter Dose Not Needed
 Starter Dose: 100 mg SQ at Week 0 and Week 4 QTY: _____ 1 Pen / 1 Syringe | Refills: 0
 Maintenance Dose: inject 80 mg SQ once every 4 weeks (starting at Week 4) QTY: _____ 1 Pen / 1 Syringe | Refills: _____

Xeljanz® 5 mg Tablet
 5 mg PO BID QTY: _____ | Refills: _____

Xeljanz® XR 11 mg Tablet
 11 mg PO once daily QTY: _____ | Refills: _____

Other: _____ QTY: _____ | Refills: _____

Prescriber's Signature: _____ DAW (Dispense as Written) **Date:** _____

Prescriber certifies that this referral form contains an original signature and is signed by the treating prescriber. NO STAMPED SIGNATURES WILL BE ACCEPTED. Where required by law, send electronic prescription or on official state prescription blank. In the event requested agent is not available through AcariaHealth, this prescription shall be forwarded to an eligible pharmacy.

IMPORTANT NOTICE: This message may contain privileged and confidential information and is intended only for the individual named. If you are not the named addressee, you should not disseminate, distribute or copy this fax. Please notify the sender immediately if you have received this document by mistake, then destroy this document. Please direct all verification or notification to AcariaHealth or any of its subsidiaries using the contact information provided on this coversheet.