

Phone: 866.892.1580 • Fax: 866.892.2363

RHEUMATOLOGY IV ROUTE REFERRAL FORM

PATIENT INFORMATION			
Patient Name:		DOB:	Sex: <input type="checkbox"/> M <input type="checkbox"/> F
SSN:		Phone:	Weight: <input type="checkbox"/> lbs. <input type="checkbox"/> kg.
Address:		Allergies:	
Emergency Contact:		City:	State: _____ Zip: _____
Phone:		<input type="checkbox"/> Please attach demographic information	
PRESCRIBER INFORMATION			
Prescriber:		NPI:	DEA: _____ State Lic: _____
Supervising Physician:		Practice Name:	
Address:		City:	State: _____ Zip: _____
Phone:		Fax:	Key Office Contact: _____ Phone: _____
DIAGNOSIS INFORMATION / MEDICAL ASSESSMENT			
Primary Diagnosis: <input type="checkbox"/> M06.9 Rheumatoid Arthritis <input type="checkbox"/> L40.54; L40.59 Psoriatic Arthritis <input type="checkbox"/> M08.00 Polyarticular Juvenile Rheumatoid Arthritis <input type="checkbox"/> M08.00 Juvenile Idiopathic Arthritis <input type="checkbox"/> M45.9 Ankylosing Spondylitis <input type="checkbox"/> M45.9 Ankylosing Spondylitis <input type="checkbox"/> M33.20 Polmyositis <input type="checkbox"/> M15.0 - M15.9 Osteoarthritis <input type="checkbox"/> Other: _____			
<input type="checkbox"/> Has patient been treated <i>previously</i> for this condition? <input type="checkbox"/> Yes <input type="checkbox"/> No Is patient <i>currently</i> on therapy? <input type="checkbox"/> Yes <input type="checkbox"/> No Medications: _____			
<input type="checkbox"/> Is patient currently on therapy? <input type="checkbox"/> Yes <input type="checkbox"/> No If yes, how long should patient wait before starting the new medication? _____			
<input type="checkbox"/> Will patient stop taking the above medication(s) before starting the new medication? <input type="checkbox"/> Yes <input type="checkbox"/> No If yes, how long should patient wait before starting the new medication? _____			
<input type="checkbox"/> Other medications patient is currently taking including OTC medications with dosage and direction (or fax medication profile): _____			
<input type="checkbox"/> Has patient received a Quatiferon gold , Tspot , or PPD (tuberculosis) Skin Test ? <input type="checkbox"/> Yes <input type="checkbox"/> No Date: _____ Results: <input type="checkbox"/> Negative <input type="checkbox"/> Positive <i>Prior to initiating treatment and periodically during therapy, patient should be evaluated for active tuberculosis and tested for latent infection</i>			
INSURANCE INFORMATION			
<input type="checkbox"/> Please attach front and back of patient's insurance card (medical and prescription)			
COPAY CARD ENROLLMENT			
<input type="checkbox"/> Please check if enrolling in copay card		Copay ID: _____	
PRESCRIPTION INFORMATION			
<input type="checkbox"/> Actemra® 20 mg/mL (Vial Sizes: 4 mL, 10 mL, 20 mL) *Maximum dose per infusion is 800 mg <input type="checkbox"/> Starter Dose: 4 mg/kg _____ mg IV every 4 weeks, infusion over 60 minutes (infusion at MD's office or infusion center) <input type="checkbox"/> Maintenance Dose: 8 mg/kg _____ mg IV every 4 weeks, infusion over 60 minutes (infusion at MD's office or infusion center) <input type="checkbox"/> Alternate Dose: _____		<input type="checkbox"/> Enroll in Access Solutions QTY: <u>QS</u> Refills: _____ QTY: <u>QS</u> Refills: _____	
<input type="checkbox"/> Epipen® 0.3 mg IM x 1, may repeat #2 <input type="checkbox"/> Epipen® Jr. for Pediatrics less than 30 kg, 0.15 mg IM x 1, may repeat <small>*In the case of anaphylaxis, inj. in anterolateral thigh area #2</small>		QTY: _____ Refills: _____	
<input type="checkbox"/> IVIG <input type="checkbox"/> Home Infusion (please go to www.AcariaHealth.com to download IVIG referral form, or ask your local AcariaHealth representative) <input type="checkbox"/> MD's office infusion <input type="checkbox"/> Infusion supplies needed <input type="checkbox"/> Dosage: _____		QTY: _____ Refills: _____ QTY: <u>QS</u> Refills: _____ QTY: <u>QS</u> Refills: _____	
<input type="checkbox"/> Kevzara® 200 mg/1.14mL Prefilled Syringe, 200 mg SQ every 2 weeks		QTY: <u>1 box (2 syringes)</u> Refills: _____	
<input type="checkbox"/> Orencia® IV 250 mg Vial for IV *Dosage based on patient's weight <input type="checkbox"/> MD's office infusion <input type="checkbox"/> Infusion supplies needed <input type="checkbox"/> Administer 125mg by subcutaneous injection once weekly with or without an intravenous loading dose. <input type="checkbox"/> For patients initiating therapy with an IV loading dose, administer a single IV infusion (per Kg categories above), followed by the first 125 mg subcutaneous injection given within a day of the IV infusion		<input type="checkbox"/> Enroll in The Circle Program QTY: _____ Refills: _____ QTY: _____ Refills: _____ QTY: _____ Refills: _____	
<input type="checkbox"/> Orencia® ClickJet Autoinjector <input type="checkbox"/> Administer 125mg by subcutaneous injection once weekly with or without an intravenous loading dose. <input type="checkbox"/> For patients initiating therapy with an IV loading dose, administer a single IV infusion (per Kg categories above), followed by the first 125 mg subcutaneous injection given within a day of the IV infusion		QTY: _____ Refills: _____	
<input type="checkbox"/> Remicade® 100 mg Vial <input type="checkbox"/> MD's office infusion <input type="checkbox"/> Infusion supplies needed <input type="checkbox"/> Starter Dose: _____ mg/kg _____ mg IV on: Week 0, 2, 6, then follow maintenance dosing instructions <input type="checkbox"/> Maintenance Dose: _____ mg/kg _____ mg IV every _____ weeks for _____ infusions		<input type="checkbox"/> Enroll in AccessOne Program QTY: _____ Refills: _____ QTY: _____ Refills: _____	
<input type="checkbox"/> Rituxan® 1000 mg IV Infusion as Directed (MD's Office Infusion) <input type="checkbox"/> Day 1 <input type="checkbox"/> Day 15 (will dispense available vial size) <input type="checkbox"/> Other: _____		<input type="checkbox"/> Enroll in RISE Program QTY: <u>QS</u> Refills: _____	
<input type="checkbox"/> Simponi® Aria 50 mg/4 mL Single-use Vial <input type="checkbox"/> Starting Dose: 2 mg/kg _____ mg IV at Weeks 0 and 4, infusion over 30 minutes <input type="checkbox"/> Maintenance Dose: 2 mg/kg _____ mg IV every 8 weeks, infusion over 30 minutes <input type="checkbox"/> Alternate Dose: _____		<input type="checkbox"/> Enroll in RISE Program QTY: <u>QS</u> Refills: _____ QTY: <u>QS</u> Refills: _____ QTY: <u>QS</u> Refills: _____	
<input type="checkbox"/> Other <input type="checkbox"/> Dosage: _____		QTY: _____ Refills: _____	

Prescriber's Signature: _____ DAW (Dispense as Written) **Date:** _____

Prescriber certifies that this referral form contains an original signature and is signed by the treating prescriber. NO STAMPED SIGNATURES WILL BE ACCEPTED. Where required by law, send electronic prescription or on official state prescription blank. In the event requested agent is not available through AcariaHealth, this prescription shall be forwarded to an eligible pharmacy.

IMPORTANT NOTICE: This message may contain privileged and confidential information and is intended only for the individual named. If you are not the named addressee, you should not disseminate, distribute or copy this fax. Please notify the sender immediately if you have received this document by mistake, then destroy this document. Please direct all verification or notification to AcariaHealth or any of its subsidiaries using the contact information provided on this coversheet.